

Limits on violations of Lorentz symmetry and the Einstein equivalence principle using radio-frequency spectroscopy of atomic dysprosium

M. A. Hohensee,* N. Leeper, and D. Budker

Physics Department, University of California, Berkeley 94720, USA

C. Harabati, V. A. Dzuba, and V. V. Flambaum

School of Physics, University of New South Wales, Sydney 2052, Australia

(Dated: March 19, 2013)

We report a joint test of local Lorentz invariance and the Einstein Equivalence Principle for electrons, using long-term measurements of the transition frequency between two nearly degenerate states of atomic dysprosium. We present many-body calculations which demonstrate that the energy splitting of these states is particularly sensitive to violations of both special and general relativity. We limit Lorentz violation for electrons at the level of 10^{-17} , matching or improving the best laboratory and astrophysical limits by up to a factor of 10, and gravitational redshift anomalies for electrons by two orders of magnitude, to 10^{-8} . With some enhancements, our experiment may be sensitive to Lorentz violation at the level of 9×10^{-20} .

PACS numbers: 03.30.+p, 04.80.-y, 11.30.Cp, 32.30.Jc

Local Lorentz invariance (LLI) and the Einstein equivalence principle (EEP) are fundamental to both the standard model and general relativity [1]. Nevertheless, these symmetries may be violated at experimentally accessible energy scales due to spontaneous symmetry breaking, or some other mechanism at high energy scales [2, 3]. This has motivated the development of many experimental tests of both LLI and EEP [4, 5], and of a phenomenological framework, known as the standard model extension (SME), which can be used to quantitatively compare these tests' results to one another [6]. This widely used [4] framework augments the standard model Lagrangian with every combination of standard model fields that are not term-by-term invariant under Lorentz transformations, while maintaining gauge invariance, energy-momentum conservation, and Lorentz invariance of the total action [6]. Violations of LLI, which themselves constitute violations of EEP [1, 5], have also been shown to violate other tenets of general relativity [7].

In this letter, we show, using many-body calculations, that the energies of two low-lying excited states of dysprosium (Dy) [8–11] are extremely sensitive to physics that breaks LLI and the EEP in the dynamics of electrons. We report the results of an analysis of Dy spectroscopy data acquired over two years that significantly improves upon the best laboratory [12] and accelerator [13] limits on electron violations of LLI and EEP. Our result is competitive with some astrophysical bounds [14]. We also improve constraints on electron-related gravitational redshift anomalies [15] by two orders of magnitude [16].

The EEP and LLI require that spacetime, while it may be curved, be locally flat, and Lorentzian [1]. Thus the relative frequencies of any set of clocks at relative rest

and located at the same point in (or within a sufficiently small volume of) spacetime be independent of a) where that point is located in a gravitational potential, or b) the velocity of their rest frame (LLI). In the SME, violation of EEP and LLI for electrons can be described by modifying the electron dispersion relation, which in turn causes the energies of bound electronic states to vary with the velocity and gravitational potential of their rest frame [7, 17].

We focus on the symmetric, traceless $c_{\mu\nu}$ tensor in the electron sector of the SME, written using coordinates such that the speed of light is a constant c in all frames. The $c_{\mu\nu}$ tensor modifies the kinetic term in the electronic QED Lagrangian to become [6]

$$\mathcal{L} = \frac{1}{2}i\bar{\psi}(\gamma_\nu + c_{\mu\nu}\gamma^\mu)\overleftrightarrow{D}^\nu\psi - \bar{\psi}m\psi, \quad (1)$$

where m is the electron mass, ψ is a four-component Dirac spinor, γ^ν are the Dirac gamma matrixes, and $f\overleftrightarrow{D}^\nu g \equiv fD^\nu g - gD^\nu f$, with $D^\nu \equiv \partial^\nu - qA^\nu$. The $c_{\mu\nu}$ tensor is frame-dependent [6, 17–19], and is uniquely specified by its value in a standard reference frame. We use the Sun-centered, celestial equatorial frame (SCCEF) for this purpose, indicated by the coordinate indexes (T, X, Y, Z) , for ease of comparison with other results [4]. The component indexes for laboratory frame coordinates are given as $(0, 1, 2, 3)$, where $t = x_0/c$ is the time coordinate. Roman indexes are used to indicate the spatial components of $c_{\mu\nu}$, and are capitalized in the SCCEF frame. The $c_{\mu\nu}$ tensor has six parity-even components: c_{TT} , plus the five c_{JK} 's; and three parity-odd components: c_{TJ} , which introduce direction and frame dependent anisotropies in the electrons' energy-momentum, or dispersion relation [6]. This shifts the energies of bound electronic states as a function of the states' orientation and alignment in absolute space, breaking both LLI and rotational symmetry [17]. In a gravitational potential, Eq. (1) acquires

* hohensee@berkeley.edu




FIG. 1. Energy levels of Dysprosium. Atoms are optically pumped (solid lines) to a state which decays (wavy lines) into the metastable state B . A linearly polarized rf field drives the $B \rightarrow A$ transition, which is detected via fluorescence at 564nm. Insets a) and b) show the magnified diagram for ^{164}Dy and ^{162}Dy , respectively. Lorentz-symmetry violation shifts the rf resonance by $\delta\omega_{\text{rf}} = (\Delta E_B - \Delta E_A)/\hbar$. The sign of the observed shift depends on the sign of the level splitting.

additional terms proportional to $c_{\mu\nu}$ and to the curvature of spacetime [7]. These arise due to the interplay between the LLI-preserving distortion of spacetime due to gravity, and the LLI-violating distortion due to $c_{\mu\nu}$, generating anomalous gravitational redshifts that scale with the electron's kinetic energy [7, 16].

In terms of spherical-tensor operators, Eq. (1) produces a shift δh in the effective Hamiltonian for a bound electron with momentum $\vec{p}$ given by [7, 17, 20]

$$\delta h = - \left(C_0^{(0)} - \frac{2U}{3c^2} c_{00} \right) \frac{\vec{p}^2}{2m} - \sum_{q=-2}^2 \frac{(-1)^q}{6m} C_q^{(2)} T_{-q}^{(2)}, \quad (2)$$

where we have included the leading order $(2U/3c^2)c_{00}$ gravitational redshift anomaly [7, 21] in terms of the Newtonian potential U , and

$$\begin{aligned} C_0^{(0)} &= c_{00} + \frac{2}{3} c_{jj}, & C_0^{(2)} &= (c_{jj} - 3c_{33}) \\ C_{\pm 1}^{(2)} &= \pm 3(c_{31} \pm ic_{32}), & C_{\pm 2}^{(2)} &= 3(c_{11} - c_{22} \pm ic_{12}), \end{aligned}$$

are written in terms of the laboratory-frame values of the $c_{\mu\nu}$ tensor, with summation implied over like indexes. Note that $C_0^{(2)}$ is also known as c_q in the literature [17]. The spherical tensor components of the squared momentum are written as $T_0^{(2)} = \vec{p}^2 - 3p_3^2$, $T_{\pm 1}^{(2)} = \pm p_3(p_1 \pm ip_2)$, and $T_{\pm 2}^{(2)} = (p_1^2 - p_2^2)/2 \pm ip_1 p_2$. The energy shift for a state $|J, M\rangle$ of an atom due to the perturbation (2) is the expectation value of the corresponding N electron operator. Since only tensors with $q = 0$ contribute to energy shifts of bound states, we need only calculate matrix elements for the $\vec{p}^2$ and $T_0^{(2)} = \vec{p}^2 - 3p_3^2$ operators.

TABLE I. Matrix elements of the relevant operators of Lorentz violation for the states A and B of Dy in units of the Hartree energy $E_h = (6.5 \times 10^{15} \text{ Hz})\hbar$.

	State A	State B
Term Symbol	$^3[10]$	$^7\text{H}^\circ$
	Energies (cm^{-1})	
Expt. [22]	19798	19798
Calc. [23]	19786	19770
	Matrix Element (units of E_h)	
$\langle J c\gamma^0(\gamma^j p_j - 3\gamma^3 p_3) J \rangle$	69.48	49.73
$\langle J \vec{p}^2 - 3p_3^2 J \rangle$	69.84	49.89
$\langle JM \vec{p}^2 JM \rangle$	437	422

Dysprosium, an atom with 66 protons and a partially filled f -shell, is well suited to measuring the electron $c_{\mu\nu}$ coefficients. It possesses two near-degenerate, low-lying excited states with significant momentum quadrupole moments, opposite parity, and leading configurations: $[\text{Xe}]4f^{10}5d6s$, $J = 10$ (state A) and $[\text{Xe}]4f^9 5d^2 6s$, $J = 10$ (state B), which differ by a transposition of an electron from the $4f$ to the $5d$ orbital. The energy difference between these states can be measured directly by driving an electric-dipole transition (Fig. 1) with a radio-frequency (rf) field, and should be particularly sensitive to anomalies proportional to the electrons' kinetic energy, since the $4f$ orbital lies partly within the radius of filled s , p , and d shells that screen the nuclear charge from the larger $5d$ orbital. The same properties of Dy that make measurements of the $B \rightarrow A$ transition particularly sensitive to variations in the fine structure constant, α [24–26], also make them highly sensitive to violations of LLI and EEP.

To calculate the relevant matrix elements for these states, we use a version of the configuration interaction method optimized for atoms with many electrons in open shells. This method has been used to calculate energy levels, transition amplitudes, dynamic polarizabilities, “magic” frequencies in optical traps, and the effects of α variation and parity violation in Dy and other atoms [27, 28]. Calculated values of the reduced matrix elements for the A and B states of Dy are presented in Table I, and details of their derivation can be found in the Supplement [29]. To check our results, we calculate the fully relativistic matrix elements of $c\gamma^0\gamma^j p_j$ and $T_0^{(2)} = c\gamma^0(\gamma^j p_j - 3\gamma^3 p_3)$, corresponding to $\vec{p}^2$ and $\vec{p}^2 - 3p_3^2$ [20]. We find good agreement between both calculations, consistent with our initial approximation.

An effusive atomic beam of Dysprosium atoms is produced by a ~ 1400 K oven, and is optically pumped into the metastable state B via consecutive laser excitations with 833 nm and 669 nm light, followed by a spontaneous decay. The atoms are resonantly excited from state B to A via an rf electric field, whose linear polarization defines the atoms' quantization axis. The polarization of the excitation laser is chosen to create a symmetric population among the $\pm M$ magnetic sublevels of state B . The A state relaxes to the ground state in a cascade




FIG. 2. Full record of frequency measurements for ^{162}Dy (upper data set) and ^{164}Dy (lower data set). Frequencies are plotted relative to 234 661 065 Hz for ^{162}Dy and 753 513 695 Hz for ^{164}Dy . Error bars are obtained by binning measurements into sets of 20 and calculating the standard error of the mean for each set. The solid line indicates the least-squares fit. Inset: an expanded view of the most recent measurements beginning on Oct. 19, 2012, with time given in Pacific Standard Time (Coordinated Universal Time minus 8 hours).

decay, emitting 564 nm light in the process. The transition frequency is determined by measuring the intensity of the 564 nm fluorescence (with a photomultiplier) as a function of radio frequency. More details regarding the experimental procedure and apparatus can be found in Refs. [25, 30].

We measure the average frequency shift of all populated magnetic sublevels of state B relative to those of state A that are coupled by the rf electric field. The energy shift of each transition is calculated using Eq. (2) and the calculated reduced matrix elements for each state. The actual distribution of population among the magnetic sublevels is found by resolving the Zeeman structure of the two states, and measuring the peak amplitude of each transition. These amplitudes are used as weights in a sum of the shifts of each state due to Eq. (2) to determine the average shift of the unresolved line. The average shift in the $B \rightarrow A$ transition frequency ω_{rf} is given by

$$\frac{\delta\omega_{\text{rf}}}{2\pi} = \pm(10^{14} \text{ Hz}) \left[500 \left(C_0^{(0)} - \frac{2U_{\odot}}{3c^2} c_{00} \right) + 9.1 C_0^{(2)} \right], \quad (3)$$

where $U_{\odot} = -M_{\odot}G/r_{\text{lab}}$ is the Sun's gravitational potential, and ω_{rf} is defined to be positive, producing a positive (negative) shift for ^{164}Dy (^{162}Dy). This sign difference helps reject background systematics, and is determined by that of the splitting between A and B . The sign of the second term depends on the relative magnetic sublevel populations.

The value of $C_0^{(0)}$ and $C_0^{(2)}$ in the laboratory-frame is a function of $c_{\mu\nu}$ in the SCCEF, and the orientation

TABLE II. Constraints on electron $c_{\mu\nu}$ -coefficients from spectroscopy of the rf transitions in ^{162}Dy and ^{164}Dy . We use the shorthand notation $c_{X-Y} \equiv c_{XX} - c_{YY}$, $c_{T(Y+Z)} \equiv c_{TY} \cos \eta + c_{TZ} \sin \eta$, and $c_{T(Y-Z)} \equiv c_{TY} \sin \eta - c_{TZ} \cos \eta$, where $\eta = 23.4^\circ$ is the angle between the Earth's spin and orbital axes. Bounds above the horizontal divider are obtained from 12 hours of continuous measurement, while those below the line are obtained from analysis of over two years of data, see text. Some uncertainties for the latter limits are adjusted for systematic error; the statistical uncertainty is then indicated in parenthesis. Past bounds on c_{JK} , c_{TJ} , and c_{TT} , and the purely gravitational limit on c_{TT} are from analyses reported in [12], [14], [13], and [16], respectively.

Combination	New Limit	Existing Limit
$0.10 c_{X-Y} + 0.99 c_{XZ}$	-9.0 ± 11	$-33 \pm 19 \times 10^{-17}$
$0.99 c_{X-Y} - 0.10 c_{XZ}$	3.7 ± 5.6	$-8.6 \pm 62 \times 10^{-17}$
$0.94 c_{XY} + 0.35 c_{YZ}$	-0.4 ± 2.8	$43 \pm 19 \times 10^{-17}$
$0.35 c_{XY} - 0.94 c_{YZ}$	3.2 ± 7.0	$5.3 \pm 23 \times 10^{-17}$
$0.98 c_{T(Y+Z)} - 0.19 c_{TX}$	$-1.3 \pm 18(3.3)$	$0.3 \pm 1.8 \times 10^{-15}$
$0.19 c_{T(Y+Z)} + 0.98 c_{TX}$	$7.3 \pm 6.6(2.4)$	$-1.5 \pm 5.4 \times 10^{-15}$
$c_{T(Y-Z)}$	$-38 \pm 30(5)$	$.011 \pm .028 \times 10^{-13}$
c_{TT}	-2.1 ± 6.7	$10^{-6}(2 \pm 2) \times 10^{-9}$
c_{TT} (gravitational)	$-14 \pm 28(9)$	$4600 \pm 4600 \times 10^{-9}$

and velocity of the lab. Thus any anomalous $\delta\omega_{\text{rf}}$ measured in the lab must vary in time [17]. The precise relation between $C_0^{(0)}$ and $C_0^{(2)}$ and the SCCEF value of $c_{\mu\nu}$ can be found in the Supplement [29]. The scalar c_{TT} component of $c_{\mu\nu}$ can be bounded via frame- or gravitational potential-dependent effects, as it contributes to the modulation of $C_0^{(2)}$, scaled by the Earth's orbital velocity squared $\beta_{\oplus}^2 \approx 1 \times 10^{-8}$, and to that of the larger scalar term in Eq.(3) via modulations of the laboratory in the Sun's gravitational potential, which have amplitude $\Delta U_{\odot}/c^2 = 1.7 \times 10^{-10}$.

Using repeated measurements of $\delta\omega_{\text{rf}}$ acquired over nearly two years, we obtain constraints on eight of the nine elements of $c_{\mu\nu}$. The c_{JK} coefficients are constrained using data collected over the course of 12 hours beginning on Oct. 19, 2012. For each isotope the mean value of 20 successive frequency measurements (~ 10 seconds) is assigned an error bar according to the standard error of the mean for that bin. The resulting data are fit to Eq. (3) in terms of c_{JK} in the SCCEF [29], augmented by an independent, constant frequency offset for each isotope. The short duration of this data set allows us to neglect the slow (1 yr^{-1} and 2 yr^{-1}) variations induced by the c_{TT} and c_{TJ} terms. These terms are neglected in this fit, as they are suppressed by at least one factor of $\beta_{\oplus} \sim 10^{-4}$, and existing limits [14] on these terms constrains their contributions well below our statistical sensitivity.

The c_{TJ} and c_{TT} coefficients are constrained using data collected between November 2010 and July 2012. The data are binned and assigned error bars as previously described. Since the above analysis of the 12 hour data set provides tight constraints on c_{JK} coefficients,




FIG. 3. Allan deviation from a two hour measurement of the ^{164}Dy transition frequency (7:00 to 9:00 on the Fig. 2 inset).

the second fit includes only the c_{TJ} and c_{TT} coefficients. The fit routine is the same as before, adding an independent linear slope for each isotope to account for long-term systematic drifts. The resulting fit includes a large signal for the combination $c_{T(Y-Z)} \equiv c_{TY} \sin \eta - c_{TZ} \cos \eta = (-38 \pm 5) \times 10^{-13}$, where η is the Earth's axial tilt. As such a signal is inconsistent with existing limits on $c_{T(Y-Z)}$ [13, 14], we suspect the presence of uncontrolled systematic shifts in $\delta\omega_{\text{rf}}$ with characteristic modulation frequencies near 1 day^{-1} and 2 day^{-1} , and amplitude 300 mHz. These systematics may be due in part to, *e.g.*, magnetic field fluctuations ($\sim 50 \text{ mHz}$), blackbody shifts due to changes in the temperature of the spectroscopy chamber ($\sim 60 \text{ mHz}$) [31], and changes in electronic offsets ($\sim 140 \text{ mHz}$). Daily fluctuations in these systematic shifts have less effect on our bounds on c_{TX} and $c_{T(Y+Z)} \equiv c_{TY} \cos \eta + c_{TZ} \sin \eta$, as these are primarily sensitive to the yearly modulation signal produced by the larger scalar component of Eq. (3) [29]. In the presence of correlated noise, our statistical error bars overstate our measurement's precision, thus we repeat the least-squares analysis without flipping signs for ^{162}Dy relative to ^{164}Dy . This model is insensitive to Lorentz violation, but is sensitive to systematic error. Where they are larger, the absolute mean of each term in this fit replaces the statistical error estimated by the original fit.

We have also analyzed our results as a test of the gravitational redshift for electrons in the Sun's gravitational potential by fitting the long term data to terms proportional to the gravitational potential, neglecting frame-dependent effects. We obtain a purely gravitational limit on the electron's c_{TT} coefficient of $-14 \pm 28 \times 10^{-9}$.

The data and fits are shown in Fig. 2. The fit results are displayed in Table II with uncertainties quoted for 68% confidence limits. The reduced chi-squared, $\bar{\chi}^2$, for the short and long time-scale fits are 1.2 and 1.8, respectively. The larger $\bar{\chi}^2$ of the long-term fit is likely due to uncontrolled systematics that have not been accounted for in our purely statistical estimation of error bars. To obtain conservative estimates on parameter uncertainties

we have scaled the statistical error bars in both fits to provide $\bar{\chi}^2 = 1$.

We have tightened experimental limits on four of the six parity-even components of the $c_{\mu\nu}$ tensor by factors ranging from 2 to 10 [4, 12]. We report limits on two combinations of the parity-odd c_{TJ} that are on par with those set by $\sim 50 \text{ TeV}$ astrophysical phenomena [4, 14]. We improve bounds on electron-related anomalies in the gravitational redshift by a factor of 160, to 2.8×10^{-8} . With optimization, our experiment could yield significantly improved constraints. As Fig. 3 shows, our experiment is statistically sensitive to $C_0^{(2)} = c_{jj} - 3c_{33}$ in the lab at the level of 2.2×10^{-16} after 400 seconds of averaging. At present, we must wait a full day for the Earth to rotate the laboratory in the fixed reference frame, increasing our susceptibility to systematics varying on that timescale. This could be addressed by active rotation of the entire apparatus, or of the polarization of the rf electric field, making possible statistically limited sensitivities to $C_0^{(2)}$ at the level of 1.5×10^{-17} in one day, and 7.8×10^{-19} in a year. Optically pumping the atoms to the $M = \pm 10$ states could increase the experiment's sensitivity to $C_0^{(2)}$ by a factor of ~ 4.5 . Increasing the interaction time of the atoms in the rf field could gain another factor of two, as the measured linewidth of 40 kHz is twice the natural linewidth of state A . An optimized experiment may thus reach sensitivities at the order of 8.7×10^{-20} in one year. This would be three orders of magnitude better than the presently reported limits on c_{JK} , two orders of magnitude better than the best sensitivities attainable by existing optical resonator tests [32], and could prove more sensitive than astrophysical tests [14, 33]. Still narrower linewidths are possible in spectroscopic measurements of the Zeeman and hyperfine structure of the ground state of trapped Dy [34], other rare-earth elements, and of the long-lived states of rare-earth ions in doped materials. Optical transition energies in trapped ion or neutral atom clocks, and of the electronic and ro-vibrational states of molecules may also be sensitive to $c_{\mu\nu}$. The latter might also probe violations of LLI and EEP for nuclei. The derivation of the scalar and quadrupole moments of the involved states will be the subject of future work.

We are grateful to Alan Kostelecký, Paul Hamilton, and Jay Tasson for stimulating discussions. D.B. acknowledges support from the Miller Institute for Basic Research in Science. We are grateful to Arman Cingöz, Valeriy Yashchuk, Alain Lapierre, and Tuan Nguyen for designing and building the Dy atomic-beam machine. We thank Holger Müller for support of this work. This work is supported by the Australian Research Council, the National Science Foundation, and the Foundational Questions Institute.

- [1] C.W. Misner, K.S. Thorne, and J.A. Wheeler, *Gravitation* (Freeman, San Francisco, 1970).
- [2] V.A. Kostelecký and S. Samuel, Phys. Rev. D **39**, 683 (1989).
- [3] T. Damour, Class. Quant. Grav. **13**, A33 (1996).
- [4] V.A. Kostelecký and N. Russell, Rev. Mod. Phys. **83**, 11 (2011).
- [5] C. M. Will, Living Reviews in Relativity **9**, 3 (2006).
- [6] D. Colladay and V.A. Kostelecký, Phys. Rev. D **55**, 6760 (1997); **58**, 116002 (1998); V.A. Kostelecký, Phys. Rev. D **69**, 105009 (2004).
- [7] V.A. Kostelecký and J.D. Tasson, Phys. Rev. D **83**, 016013 (2011).
- [8] V.A. Dzuba, V.V. Flambaum, I.B. Khriplovich, Z. Phys. D: *Atoms, Molecules and Clusters*, **1**, 243-245 (1986).
- [9] V.A. Dzuba, V.V. Flambaum, and J.K. Webb, Phys. Rev. Lett., **82**, 888 (1999); Phys. Rev. A, **59**, 230 (1999).
- [10] V.A. Dzuba, V.V. Flambaum, and M.G. Kozlov, Phys. Rev. A, **50**, 3812 (1994).
- [11] V.A. Dzuba, V.V. Flambaum, and M.V. Marchenko, Phys. Rev. A, **68**, 022506 (2003).
- [12] H. Müller, et al., Phys. Rev. Lett. **99**, 050401 (2007).
- [13] B. Altschul, Phys. Rev. D **82**, 016002 (2010).
- [14] B. Altschul, Phys. Rev. Lett. **96**, 201101 (2006); Phys. Rev. D **74**, 083003 (2006).
- [15] R.F.C. Vessot et al., Phys. Rev. Lett. **45**, 2081 (1980).
- [16] M.A. Hohensee, S. Chu, A. Peters, and H. Müller, Phys. Rev. Lett. **106**, 151102 (2011).
- [17] V.A. Kostelecký, C.D. Lane, Phys. Rev. D **60**, 116010 (1999).
- [18] R. Bluhm, V.A. Kostelecký, C.D. Lane, and N. Russell, Phys. Rev. Lett., **88**, 090801 (2002); Phys. Rev. D **68**, 125008 (2003).
- [19] B. Altschul, Phys. Rev. D **81**, 041701 (2010).
- [20] V.A. Kostelecký, C.D. Lane, J. Math. Phys. (N.Y.) **40**, 6245 (1999).
- [21] M.A. Hohensee and H. Müller, to be published.
- [22] W.C. Martin, R. Zalubas, and L. Hagan, *Atomic Energy Levels - The Rare-Earth Elements*, NIST, Washington (1978).
- [23] V.A. Dzuba and V.V. Flambaum, Phys. Rev. A. **81**, 052515 (2010).
- [24] A.T. Nguyen and D. Budker and S. K. Lamoreaux and J. R. Torgerson, Phys. Rev. A **69**, 022105 (2004).
- [25] A. Cingöz, A. Lapierre, A.-T. Nguyen, N. Leefer, D. Budker, S. K. Lamoreaux, and J. R. Torgerson, Phys. Rev. Lett. **98**, 040801 (2007); S.J. Ferrell, A. Cingöz, A. Lapierre, A.-T. Nguyen, N. Leefer, D. Budker, V.V. Flambaum, S.K. Lamoreaux, and J.R. Torgerson, Phys. Rev. A **76**, 062104 (2007).
- [26] N.A. Leefer *et al.*, to be published.
- [27] V.A. Dzuba and V.V. Flambaum, Phys. Rev. A, **77**, 012514 (2008); Phys. Rev. A, **77**, 012515 (2008).
- [28] V.A. Dzuba, Phys. Rev. A, **71**, 032512 (2005); V.A. Dzuba and V.V. Flambaum, Phys. Rev. A. **75**, 052504 (2007).
- [29] See Supplemental Material that follows for the derivation of the matrix elements of the states *A* and *B*, and the expression for Eq. (3) in terms of $c_{\mu\nu}$ in the SCCEF.
- [30] N. Leefer, A. Cingöz, D. Budker, S.J. Ferrell, V.V. Yashchuk, A. Lapierre, A.-T. Nguyen, S.K. Lamoreaux,

and J.R. Torgerson, in *Proceedings of the 7th Symposium Frequency Standards and Metrology, Asilomar, October 2008*, edited by Lute Maleki, World Scientific, pp. 34-43.

- [31] N.A. Leefer *et al.*, to be published.
- [32] S. Herrmann and A. Senger and K. Möhle and M. Nagel and E.V. Kovalchuk and A. Peters, Phys. Rev. D. **80**, 105011 (2009); Ch. Eisele, A.Yu. Nevsky and S. Schiller, Phys. Rev. Lett. **103**, 090401 (2009).
- [33] F.R. Klinkhamer and M. Schreck, Phys. Rev. D **78**, 085026 (2008).
- [34] M. Lu, S.H. Youn, and B.L. Lev, Phys.. Rev. Lett. **104**, 063001 (2010).
- [35] R. Szymkowski, J. Math. Chem. **42**, 397 (2007).

SUPPLEMENTARY MATERIALS: MATRIX ELEMENTS

Here, we present the derivation of the explicit form of the single orbital matrix elements for the operators $\vec{p}^2$ and $T_0^{(2)} = \vec{p}^2 - 3p_3^2$. We also present the matrix elements for the relativistic operators $c\gamma^0\gamma^j p_j$ and $T_0^{(2)} = c\gamma^0(\gamma^j p_j - 3\gamma^3 p_3)$.

The many-electron state of an atom is a linear combination of Slater determinants, constructed from single-electron orbitals. These orbitals, in the relativistic limit, can be written as

$$\varphi_{n\kappa m}(\vec{r}) = \frac{1}{r} \begin{pmatrix} f_{n\kappa} \Omega_{\kappa m}(\theta, \phi) \\ i\alpha g_{n\kappa}(r) \Omega_{-\kappa m}(\theta, \phi) \end{pmatrix}, \quad (4)$$

where $\kappa = \mp(j + 1/2)$ (with $j = l \pm 1/2$) is the quantum number for the angular momentum of a Dirac spinor, α is the fine structure constant, and the spin-dependent component $\Omega_{\kappa m}(\theta, \phi)$ is given by

$$\Omega_{\kappa m}(\theta, \phi) = \begin{pmatrix} \pm \sqrt{\frac{\kappa+1/2-m}{2\kappa+1}} Y_{l,m-1/2}(\theta, \phi) \\ \sqrt{\frac{\kappa+1/2+m}{2\kappa+1}} Y_{l,m+1/2}(\theta, \phi) \end{pmatrix}, \quad (5)$$

where $Y_{l,m}(\theta, \phi)$ is a spherical harmonic. The upper component of Eq. (4) corresponds to the electron component of the spinor wave function, while the lower component to the positron. In the non-relativistic limit, we can drop the antimatter component of Eq. (4) and evaluate matrix elements in terms of

$$\varphi_{n\kappa m}(\vec{r}) = \frac{f_{n\kappa}(r)}{r} \Omega_{\kappa m}(\theta, \phi). \quad (6)$$

Using the Pauli matrix identity $(\vec{\sigma} \cdot \vec{p})^2 = \vec{p}^2$, with $\vec{\sigma}$ a vector of Pauli matrixes, we can write the action of the operator $\vec{p}^2$ on $\varphi_{n\kappa m}(\vec{r})$ as

$$(\vec{\sigma} \cdot \vec{p})^2 \varphi_{n\kappa m}(\vec{r}) = -\frac{\hbar^2}{r} \left(\frac{d^2 f_{n\kappa}}{dr^2} - \frac{\kappa(\kappa+1)}{r^2} f_{n\kappa} \right) \Omega_{\kappa m}. \quad (7)$$

The matrix elements for $\vec{p}^2$ are thus equal to $\delta_{\kappa'\kappa}I_1$, where I_1 is the radial integral

$$I_1 = \hbar^2 \int_0^\infty dr \left(\frac{\partial f_{n'\kappa'}}{\partial r} \frac{\partial f_{n\kappa}}{\partial r} + \frac{\kappa(\kappa+1)}{r^2} f_{n'\kappa'} f_{n\kappa} \right). \quad (8)$$

Similarly, the matrix elements of $p_3^2 = -\hbar^2 \partial^2 / \partial z^2$ can be found by applying $p_3 = -i\hbar \partial / \partial z$ to the spinor twice (*e.g.*, by using Eq. (3.3.3) of [35]). Using the Wigner-Eckart theorem, the matrix elements of $T_0^{(2)}$ can be written in terms of reduced matrix elements after factoring out the $3j$ -symbol:

$$\langle n'\kappa'm | T_0^{(2)} | n\kappa m \rangle = (-1)^{j'-m} \begin{pmatrix} j' & 2 & j \\ -m & 0 & m \end{pmatrix} \times \langle n'\kappa' || T^{(2)} || n\kappa \rangle. \quad (9)$$

These reduced matrix elements are then given by

$$\langle n'\kappa' || T^{(2)} || n\kappa \rangle = \begin{cases} A(j', j)I_1 & \text{if } \kappa' = -\kappa - 1 \\ A(j', j)I_2 & \text{if } \kappa' = -\kappa + 1 \\ B(j', j)I_2 & \text{if } \kappa' = \kappa - 2 \\ B(j', j)I_3 & \text{if } \kappa' = \kappa + 2 \\ C(j)I_1 & \text{if } \kappa' = \kappa \end{cases}, \quad (10)$$

where $A(j', j)$, $B(j', j)$, and $C(j)$ are given by

$$A(j', j) = (-1)^{j_m - j' + 1} \left[\frac{6(2j_m + 3)(2j_m + 1)}{(2j_m + 2)2j_m(2j_m + 4)} \right]^{1/2}$$

$$B(j', j) = (-1)^{j_m - j' + 1} \left[\frac{3(2j_m + 5)(2j_m + 3)(2j_m + 1)}{2(2j_m + 4)(2j_m + 2)} \right]^{1/2}$$

$$C(j) = \left[\frac{(2j + 3)(2j + 1)(2j - 1)}{4j(j + 1)} \right]^{1/2}$$

where $j_m = \min(j', j)$. The radial integrals I_2 and I_3 are

$$I_2 = \hbar^2 \int_0^\infty dr \left(\frac{\partial f_{n'\kappa'}}{\partial r} \frac{\partial f_{n\kappa}}{\partial r} - \frac{2\kappa - 1}{r} f_{n'\kappa'} \frac{\partial f_{n\kappa}}{\partial r} - \frac{\kappa(\kappa - 2)}{r^2} f_{n'\kappa'} f_{n\kappa} \right)$$

and

$$I_3 = \hbar^2 \int_0^\infty dr \left(\frac{\partial f_{n'\kappa'}}{\partial r} \frac{\partial f_{n\kappa}}{\partial r} + \frac{2\kappa + 3}{r} f_{n'\kappa'} \frac{\partial f_{n\kappa}}{\partial r} - \frac{(\kappa + 3)(\kappa + 1)}{r^2} f_{n'\kappa'} f_{n\kappa} \right).$$

As for the the matrix elements of the relativistic form of the operators acting on the four-component spinor in Eq. (4), we find that their angular components $A(j', j)$,

$B(j', j)$, and $C(j)$ are the same, while the radial integrals differ, so that

$$\langle n'\kappa' || c\gamma^0(\gamma^j p_j - 3\gamma^3 p_3) || n\kappa \rangle = \begin{cases} A(j', j)\tilde{I}_1 & \text{if } \kappa' = -\kappa - 1 \\ A(j', j)\tilde{I}_2 & \text{if } \kappa' = -\kappa + 1 \\ B(j', j)\tilde{I}_3 & \text{if } \kappa' = \kappa - 2 \\ B(j', j)\tilde{I}_4 & \text{if } \kappa' = \kappa + 2 \\ C(j)\tilde{I}_5 & \text{if } \kappa' = \kappa \end{cases}, \quad (11)$$

where

$$\begin{aligned} \tilde{I}_1 &= -\frac{c\alpha\hbar}{2} \int_0^\infty dr \left((2\kappa - 1)g_{n'\kappa'} \frac{\partial f_{n\kappa}}{\partial r} + (2\kappa + 3)f_{n'\kappa'} \frac{\partial g_{n\kappa}}{\partial r} - \frac{(2\kappa - 1)(\kappa + 1)}{r} g_{n'\kappa'} f_{n\kappa} - \frac{(2\kappa + 3)\kappa}{r} f_{n'\kappa'} g_{n\kappa} \right), \\ \tilde{I}_2 &= -\frac{c\alpha\hbar}{2} \int_0^\infty dr \left((2\kappa - 3)g_{n'\kappa'} \frac{\partial f_{n\kappa}}{\partial r} + (2\kappa + 1)f_{n'\kappa'} \frac{\partial g_{n\kappa}}{\partial r} + \frac{(2\kappa - 3)\kappa}{r} g_{n'\kappa'} f_{n\kappa} + \frac{(2\kappa + 1)(\kappa - 1)}{r} f_{n'\kappa'} g_{n\kappa} \right), \\ \tilde{I}_3 &= -2c\alpha\hbar \int_0^\infty dr \left(f_{n'\kappa'} \frac{\partial g_{n\kappa}}{\partial r} + \frac{\kappa - 1}{r} f_{n'\kappa'} g_{n\kappa} \right), \\ \tilde{I}_4 &= 2c\alpha\hbar \int_0^\infty dr \left(g_{n'\kappa'} \frac{\partial f_{n\kappa}}{\partial r} - \frac{\kappa + 1}{r} g_{n'\kappa'} f_{n\kappa} \right), \\ \tilde{I}_5 &= c\alpha\hbar \int_0^\infty dr \left(g_{n'\kappa'} \frac{\partial f_{n\kappa}}{\partial r} - f_{n'\kappa'} \frac{\partial g_{n\kappa}}{\partial r} + \frac{\kappa}{r} g_{n'\kappa'} f_{n\kappa} + \frac{\kappa}{r} f_{n'\kappa'} g_{n\kappa} \right). \end{aligned}$$

SUPPLEMENTARY MATERIALS: MANY-BODY CALCULATION

To evaluate the expectation of the electronic kinetic energy for states A and B in Dy, we use a version of the configuration interaction (CI) method, originally developed for calculating energy levels, electromagnetic amplitudes, dynamics polarizabilities, “magic” frequencies in optical traps, the effects of variations in the fine structure constant, and parity violation in atoms with many electrons in open shells [27]. The effective Hamiltonian for N_v valence electrons in an atom ($N_v = 12$ for Dy) has the form

$$\hat{H}^{\text{eff}} = \sum_{j=1}^{N_v} \hat{h}_{1j}(r_j) + \sum_{j < k}^{N_v} e^2 / r_{jk}, \quad (12)$$

where e is the elementary electron charge, and $\hat{h}_{1j}$ is the one-electron Hamiltonian

$$\hat{h}_{1j}(r_j) = c\vec{\alpha} \cdot \vec{p} + (\beta - 1)m_e c^2 - \frac{Ze^2}{r} + V_{\text{core}} + \delta V, \quad (13)$$

where $\alpha_j = \gamma^0 \gamma_j$, and $\beta = \gamma^0$. Here, V_{core} is the Hartree-Fock potential due to the core electrons, and δV simulates the effects of correlations between core and valence

TABLE III. Configurations and effective core polarizabilities α_p (a.u.) used in the calculations.

N	Parity	Configuration	α_p
1	Even	$4f^{10}6s^2$	0.4
2	Even	$4f^{10}6s5d$	0.4006
3	Even	$4f^96s^26p$	0.4039
4	Even	$4f^95d6s6p$	0.389
5	Even	$4f^{10}6p^2$	0.4
6	Even	$4f^95d^26p$	0.4
7	Odd	$4f^95d^26s$	0.3947
8	Odd	$4f^95d6s^2$	0.3994
9	Odd	$4f^{10}6s6p$	0.397
10	Odd	$4f^{10}5d6p$	0.4
11	Odd	$4f^95d6p^2$	0.4
12	Odd	$4f^96s6p^2$	0.4

electrons. It is also known as the polarization potential, and has the form

$$\delta V = -\frac{\alpha_p}{2(r^4 + a^4)}, \quad (14)$$

where α_p is the core polarizability, and a is a cutoff parameter (here, the Bohr radius $a = a_B$). Table III lists the configurations considered in our calculation. The self-consistent Hartree-Fock procedure is performed separately for each configuration. Next, valence states obtained from the Hartree-Fock calculations are used as basis states for the CI calculation. The CI method requires that the atomic core remain the same for each outer electron configuration. We select the core state that corresponds to the ground state configuration. Changes in the core state due to changes in the valence state are small, and can be neglected, as the $6s$, $6p$ and $5d$ shells are comparatively distant, and the potential they produce at the core is a nearly uniform perturbation. We can thus model the valence potential as shifting the energy of the core eigenstates, without modifying the wave functions. The $4f$ electrons, on the other hand, lie nearer to the core, and thus have a stronger effect. In all cases considered here, however, only one of the ten $4f$ electrons

ever change state. Thus their total effect on the atomic core will also be small. A more detailed discussion of the effects of valence electrons on atomic core states can be found in Refs. [28].

The form of δV in Eq. (14) is chosen to coincide with the standard polarization potential at large distances ($-\alpha_p/2r^4$). We treat the α_p for each configuration as a fit parameter, chosen to match experimentally measured energy intervals between states of different configurations. Their values are displayed in Table III. The values of α_p are similar for all configurations considered. This is expected, as the state of the core electrons is the same for each valence configuration. The small differences in α_p for different configurations help compensate the effect of the incompleteness of our valence electron basis, and other imperfections in the calculation.

SUPPLEMENTARY MATERIALS: FRAME DEPENDENCE OF $\delta\omega_{\text{rf}}$

The laboratory frame components of the $c_{\mu\nu}$ tensor are found in terms of the sun-centered equatorial frame (SCCEF) components by requiring that the Lagrangian of Eq. (1) be invariant under the observer Lorentz transformation between frames. This condition leads to

$$c_{\mu\nu} = c_{MN}\Lambda_\mu^M\Lambda_\nu^N, \quad (15)$$

where c_{MN} is the c tensor defined in the SCCEF. In the SCCEF the Z axis points along the Earth's rotation axis, the $X-Y$ plane is the Earth's equatorial plane, and the Y axis points from the Sun to the Earth at the vernal equinox. The Lorentz transformation Λ_μ^M is a rotation, dependent on experimental geometry and the rotation of the Earth, to align the laboratory defined axes with these SCCEF axes followed by a boost, determined mainly by the Earth's orbital velocity, to the rest frame of the Sun. Our laboratory frame is defined such that z is parallel to the quantization axis defined by the polarization of the rf field. Thus z points $\theta = 15^\circ$ south of west, x points $\theta = 15^\circ$ east of south, and y points vertically upwards. The rotation from the SCCEF to the laboratory frame is given by

$$R = \begin{pmatrix} \cos\chi \cos\theta \cos\omega T_\oplus - \sin\theta \sin\omega T_\oplus & \sin\theta \cos\omega T_\oplus + \cos\chi \cos\theta \sin\omega T_\oplus & -\sin\chi \cos\theta \\ -\sin\chi \cos\omega T_\oplus & -\sin\chi \sin\omega T_\oplus & -\cos\chi \\ -\cos\chi \sin\theta \cos\omega T_\oplus - \cos\theta \sin\omega T_\oplus & \cos\theta \cos\omega T_\oplus - \cos\chi \sin\theta \sin\omega T_\oplus & \sin\chi \sin\theta \end{pmatrix}, \quad (16)$$

where $\chi = 52.1^\circ$ is the colatitude of our laboratory, ω is the angular frequency of the Earth's rotation in the SCCEF (*i.e.* $2\pi \times$ an inverse sidereal day), and $T_\oplus$ is measured from the first time that East, as measured in the laboratory, and the SCCEF Y -axis coincides after a vernal equinox. The boost of the laboratory in the

SCCEF frame is given by [4]

$$\vec{\beta} = \begin{pmatrix} \beta_\oplus \sin\Omega T - \beta_L \sin\chi \sin\omega T \\ -\beta_\oplus \cos\eta \cos\Omega T + \beta_L \sin\chi \cos\omega T \\ -\beta_\oplus \sin\eta \cos\Omega T \end{pmatrix}, \quad (17)$$

where $\eta = 23.4^\circ$ is the angle between the ecliptic plane and the Earth's equatorial plane, $\beta_\oplus =$

TABLE IV. Dominant time-varying terms in the fit for the c_{JK} coefficients. The frequencies $\omega_\oplus$ and Ω and the sidereal-day and yearly frequencies, respectively. The colatitude of the experiment is given by $\chi \sim 52.1^\circ$, $\theta \sim 15^\circ$ is the angle the quantization axis is rotated towards the South from West, and $\eta \sim 23.4^\circ$ is the angle between the ecliptic and the Earth's equatorial plane. The orbital boost is $\beta_\oplus \sim 10^{-4}$. The constants $\mathcal{S} \sim \mp 5 \times 10^{16}$ Hz and $\mathcal{Q} \sim \mp 9.1 \times 10^{14}$ Hz are the scalar and quadrupole shifts, respectively, from Eq. (3). For ease of comparison with Table II, we have defined $c_{X-Y} \equiv c_{XX} - c_{YY}$ and $c_{X+Y} \equiv c_{XX} + c_{YY}$. Additional terms of $O(\mathcal{Q}\beta_\oplus^2)$ have been suppressed, although they are included in our fits.

ω_j	S_j	C_j
$\omega_\oplus$	$3\mathcal{Q} (c_{XZ} \sin \chi \sin 2\theta + c_{YZ} \sin 2\chi \sin^2 \theta)$	$3\mathcal{Q} (c_{XZ} \sin 2\chi \sin^2 \theta - c_{YZ} \sin \chi \sin 2\theta)$
$2\omega_\oplus$	$-\frac{3}{2}\mathcal{Q} [c_{X-Y} \cos \chi \sin 2\theta - \frac{1}{2}c_{XY} (1 + 3 \cos 2\theta - 2 \cos 2\chi \sin^2 \theta)]$	$3\mathcal{Q} \left[\frac{c_{XY} \cos \chi \sin 2\theta +}{\frac{1}{8}c_{X-Y} (1 + 3 \cos 2\theta - 2 \cos 2\chi \sin^2 \theta)} \right]$
2Ω	-	$-c_{X+Y} \frac{5}{12} \mathcal{S} \beta_\oplus^2 \sin^2 \eta$

TABLE V. Dominant time-varying terms in the fit for the c_{TJ} and c_{TT} coefficients. The frequencies $\omega_\oplus$ and Ω and the sidereal-day and yearly frequencies, respectively. The colatitude of the experiment is given by $\chi \sim 52.1^\circ$, $\theta \sim 15^\circ$ is the angle the quantization axis is rotated towards the South from West, and $\eta \sim 23.4^\circ$ is the angle between the ecliptic and the Earth's equatorial plane. The orbital boost is $\beta_\oplus \sim 10^{-4}$. The constants $\mathcal{S} \sim \mp 5 \times 10^{16}$ Hz and $\mathcal{Q} \sim \mp 9.1 \times 10^{14}$ Hz are the scalar and quadrupole shifts, respectively, from Eq. (3). For ease of comparison with Table II, we have defined $c_{T(Y+Z)} \equiv c_{TY} \cos \eta + c_{TZ} \sin \eta$ and $c_{T(Y-Z)} \equiv c_{TY} \sin \eta - c_{TZ} \cos \eta$. Additional terms of $O(\mathcal{Q}\beta_\oplus)$ proportional to c_{TX} and $c_{T(Y+Z)}$ have been suppressed, although they are included in our fits. The gravitational terms appear at $\sin \Omega T$ and $\cos \Omega T$ due to a phase offset $\phi_\odot = 10.4^\circ$ between the oscillation of the boost vector (measured from the vernal equinox) and the oscillation of the Earth in the Solar gravitational potential (with perihelion on Jan. 3rd). The amplitude of the Earth's modulation in the Solar gravitational potential is $\Delta U/c^2 \approx 1.7 \times 10^{-10}$.

ω_j	S_j	C_j
Ω	$\frac{10}{3} \mathcal{S} \beta_\oplus c_{TX} + \frac{2\Delta U}{3c^2} \mathcal{S} c_{TT} \sin \phi_\odot$	$-\frac{10}{3} \mathcal{S} \beta_\oplus c_{T(Y+Z)} + \frac{2\Delta U}{3c^2} \mathcal{S} c_{TT} \cos \phi_\odot$
2Ω	-	$+\frac{3}{8} \mathcal{Q} \beta_\oplus c_{T(Y-Z)} \sin 2\eta (1 + 3 \cos 2\theta + 6 \cos 2\chi \sin^2 \theta)$
$\omega_\oplus - 2\Omega$	$2\mathcal{Q} \beta_\oplus^2 c_{TT} \cos^3 \frac{\eta}{2} \sin \frac{\eta}{2} \sin^2 \theta \sin 2\chi$	$+\frac{1}{8} \mathcal{Q} \beta_\oplus^2 c_{TT} \sin^2 \eta (1 + 3 \cos 2\theta + 6 \cos 2\chi \sin^2 \theta)$
$\omega_\oplus - \Omega$	$\frac{3}{2} \mathcal{Q} \beta_\oplus c_{T(Y-Z)} (\cos \eta + \cos 2\eta) \sin 2\chi \sin^2 \theta$	$-\frac{1}{2} \mathcal{Q} \beta_\oplus^2 c_{TT} \sin \eta (1 + \cos \eta) \sin \chi \sin 2\theta$
$\omega_\oplus$	$-\frac{1}{2} \mathcal{Q} \beta_\oplus^2 c_{TT} \sin 2\eta \sin 2\chi \sin^2 \theta$	$-\frac{3}{2} \mathcal{Q} \beta_\oplus c_{T(Y-Z)} (\cos \eta + \cos 2\eta) \sin \chi \sin 2\theta$
$\omega_\oplus + \Omega$	$-\frac{3}{2} \mathcal{Q} \beta_\oplus c_{T(Y-Z)} (\cos \eta - \cos 2\eta) \sin 2\chi \sin^2 \theta$	$-\frac{1}{2} \mathcal{Q} \beta_\oplus^2 c_{TT} \sin 2\eta \sin \chi \sin 2\theta$
$\omega_\oplus + 2\Omega$	$\frac{1}{4} \mathcal{Q} \beta_\oplus^2 c_{TT} (2 \sin \eta - \sin 2\eta) \sin 2\chi \sin^2 \theta$	$\frac{3}{2} \mathcal{Q} \beta_\oplus c_{T(Y-Z)} (\cos \eta - \cos 2\eta) \sin \chi \sin 2\theta$
$2\omega_\oplus - 2\Omega$	$\mathcal{Q} \beta_\oplus^2 c_{TT} \cos^4 \frac{\eta}{2} \cos \chi \sin 2\theta$	$\frac{1}{4} \mathcal{Q} \beta_\oplus^2 c_{TT} (2 \sin \eta - \sin 2\eta) \sin \chi \sin 2\theta$
$2\omega_\oplus - \Omega$	$\frac{3}{2} \mathcal{Q} \beta_\oplus c_{T(Y-Z)} \sin \eta (1 + \cos \eta) \cos \chi \sin 2\theta$	$-\frac{1}{4} \mathcal{Q} \beta_\oplus^2 c_{TT} \cos^4 \frac{\eta}{2} (1 + 3 \cos 2\theta - 2 \cos 2\chi \sin^2 \theta)$
$2\omega_\oplus$	$-\frac{1}{2} \mathcal{Q} \beta_\oplus^2 c_{TT} \sin^2 \eta \cos \chi \sin 2\theta$	$\frac{3}{8} \mathcal{Q} \beta_\oplus c_{T(Y-Z)} \sin \eta (1 + \cos \eta) (1 + 3 \cos 2\theta - 2 \cos 2\chi \sin^2 \theta)$
$2\omega_\oplus + \Omega$	$3\mathcal{Q} \beta_\oplus c_{T(Y-Z)} \sin \eta \sin^2 \frac{\eta}{2} \cos \chi \sin 2\theta$	$+\frac{1}{8} \mathcal{Q} \beta_\oplus^2 c_{TT} \sin^2 \eta (1 + 3 \cos 2\theta - 2 \cos 2\chi \sin^2 \theta)$
$2\omega_\oplus + 2\Omega$	$\mathcal{Q} \beta_\oplus^2 c_{TT} \sin^4 \frac{\eta}{2} \cos \chi \sin 2\theta$	$-\frac{3}{4} \mathcal{Q} \beta_\oplus c_{T(Y-Z)} \sin \eta \sin^2 \frac{\eta}{2} (1 + 3 \cos 2\theta - 2 \cos 2\chi \sin^2 \theta)$
		$-\frac{1}{4} \mathcal{Q} \beta_\oplus^2 c_{TT} \sin^4 \frac{\eta}{2} (1 + 3 \cos 2\theta - 2 \cos 2\chi \sin^2 \theta)$

$2\pi(1 \text{ a.u.})/c(1 \text{ yr}) \simeq 10^{-4}$ is the boost from the Earth's orbital velocity, $\beta_L = R_\oplus \omega \simeq 1.5 \times 10^{-6}$ is the boost from the rotational velocity of a point on the Earth's equator, $2\pi/\Omega$ is a sidereal year, and T is the time since the epoch, chosen to be the vernal equinox in the year 2000 [4]. The boost $\beta_L \sin \chi$, though included in our fits, is too small to make a significant contribution to our fits, and will henceforth be dropped from this discussion.

Since the transformation between frames is time-dependent, constant values of c_{MN} in the SCCEF give rise to time varying frequency shifts in the laboratory value of $c_{\mu\nu}$, and thus to time variations in $\delta\omega_{\text{rf}}$, via Eq. (3). Since c_{MN} also gives rise to an anomalous gravitational redshift, there is an additional contribution proportional to c_{TT} , such that $\delta\omega_{\text{rf}} \approx \mp(5 \times 10^{16} \text{ Hz}) \times 2\Delta U/(3c^2) \cos(\Omega T - \phi_\odot)$, as given in Eq. (3), where $\Delta U \sim 1.7 \times 10^{-10} c^2$ is the amplitude of the Earth's yearly

modulation in the solar gravitational potential due to the eccentricity of the Earth's orbit, and $\phi_\odot$ is such that $\cos(\Omega T - \phi_\odot)$ is minimized at perihelion ($\sim \text{Jan } 3$).

The c_{JK} coefficients contribute leading order energy shifts at much shorter time scales (daily) than the c_{TJ} and c_{TT} coefficients (yearly). As such, the c_{JK} coefficients are constrained using a single long data set acquired over the course of one day to minimize the influence of systematic effects acting on long time scales. To fit the c_{JK} terms, we assume that the parity-odd c_{TJ} coefficients are as constrained by astrophysical observations [14], and that the contribution of the c_{TT} term to the daily modulation of $\delta\omega_{\text{rf}}$ is negligible. Although we retain all terms up to $O(\beta_\oplus^2)$ in our fit function, the dominant terms that are relevant to our fit are

$$\delta\omega_{\text{rf}} = A + \sum_j (S_j \sin \omega_j T + C_j \cos \omega_j T), \quad (18)$$

where A is an independent offset parameter for each isotope, and the relevant frequencies ω_j , and quadratures S_j and C_j which contain the relevant components of c_{JK} , are summarized in Table IV. Since our experiment alternates between measuring $\delta\omega_{\text{rf}}$ for ^{162}Dy and ^{164}Dy , we do not directly compare $\delta\omega_{\text{rf}}$ for the two isotopes. We therefore perform a simultaneous fit of the separate functions (18) for each isotope, subject to the constraint that the values of c_{JK} must be the same for both.

The c_{TJ} and c_{TT} coefficients are constrained using data acquired at irregular intervals over two years. To fit the c_{TJ} and c_{TT} terms, we set c_{JK} to zero, and drop terms proportional to $\beta_{\oplus}^2$ that do not multiply c_{TT} , since existing bounds on $c_{X+Y} \equiv c_{XX} + c_{YY}$ are sufficient to ensure its doubly boost-suppressed contribution to $\delta\omega_{\text{rf}}$ is negligible at our level of precision [4]. The fit function is

$$\delta\omega_{\text{rf}} = A + MT + \sum_j (S_j \sin \omega_j T + C_j \cos \omega_j T), \quad (19)$$

where as before, A is an isotope-dependent offset, and the MT term is applied to remove any linear drifts. The relevant frequencies ω_j , and quadratures S_j and C_j , which contain the relevant components of c_{TJ} and c_{TT} , are summarized in Table V. We perform a joint fit on the ^{162}Dy and ^{164}Dy data as before, and note an apparently large signal for $c_{T(Y-Z)} \approx (38 \pm 5) \times 10^{-13}$, roughly 7.6 times the statistical error bar. As such a result would be inconsistent with other experiments [4], we suspect it may be due to modulated systematic errors that we cannot fully distinguish from our model function with the existing dataset. To estimate these errors, we fit the same data to a modified fit function that does not change sign

for the two Dy isotopes, thus maximizing our sensitivity to any common-mode systematics, and replace our original statistical error bars on each coefficient with their magnitudes in the fit to the modified function, if they are greater. If our original signal for $c_{T(Y-Z)}$ were not due to systematics, and thus truly due to Lorentz symmetry violation, we would expect to obtain a small value for $c_{T(Y-Z)}$ in the fit to the modified function. Instead, we observe that the mean value of $c_{T(Y-Z)}$ without isotopic sign reversal is nearly as large as it is for the fit to the Lorentz-violating model, and thus conclude that this is not evidence for violation of LLI. On the other hand, our estimates of the systematic error in our fits to c_{TX} and $c_{T(Y+Z)}$ is comparatively quite low. This is consistent with the hypothesis that our systematic background modulates with a period of one or half a day, and averages out over longer times, since our sensitivity to c_{TX} and $c_{T(Y+Z)}$ comes primarily from the once-yearly modulated scalar term in Eq. (3), while $c_{T(Y-Z)}$ is equally sensitive to both yearly and daily modulations proportional to the quadrupole term (see Table V).

In both cases, the uncorrelated combinations of coefficients reported in Table II were found by diagonalizing the covariance matrix from the least-squares fit to Eqs. (18) and (19).

All frequency measurements are made relative to a Cs frequency reference. Though in general, violation of LLI may affect the frequency of this reference, such effects can be neglected here. The relevant quadrupole term is strongly suppressed due to the symmetries of the $6s^1$ state [17], and the scalar shift is expected to be $\sim (10 \text{ GHz})c_{\mu\nu}$ [16], while as Table I shows, the splitting between the A and B states of Dy will be $\sim (100 \text{ THz})c_{\mu\nu}$ or greater.